

REVIEW

Human glycocalyx shedding: Systematic review and critical appraisal

Robert G. Hahn^{1,2}  | Vasu Patel³ | Randal O. Dull⁴

¹Research Unit, Södertälje Hospital, Södertälje, Sweden

²Karolinska Institute at Danderyds Hospital (KIDS), Stockholm, Sweden

³Department of Internal Medicine, Northwestern Medicine McHenry Hospital, McHenry, IL, USA

⁴Department of Anesthesiology, Pathology, Physiology, Surgery, University of Arizona, College of Medicine, Tucson, AZ, USA

Correspondence

Robert G. Hahn, MD, PhD, Professor of Anesthesia & Intensive Care, Research Unit, Södertälje Hospital, 152 86 Södertälje, Sweden.

Email: r.hahn@telia.com; robert.hahn@sll.se

Funding information

None.

Abstracts

Background: The number of studies measuring breakdown products of the glycocalyx in plasma has increased rapidly during the past decade. The purpose of the present systematic review was to assess the current knowledge concerning the association between plasma concentrations of glycocalyx components and structural assessment of the endothelium.

Methods: We performed a literature review of Pubmed to determine which glycocalyx components change in a wide variety of human diseases and conditions. We also searched for evidence of a relationship between plasma concentrations and the thickness of the endothelial glycocalyx layer as obtained by imaging methods.

Results: Out of 3,454 publications, we identified 228 that met our inclusion criteria. The vast majority demonstrate an increase in plasma glycocalyx products. Sepsis and trauma are most frequently studied, and comprise approximately 40 publications. They usually report 3-4-fold increased levels of glycocalyx degradation products, most commonly of syndecan-1. Surgery shows a variable picture. Cardiac surgery and transplantations are most likely to involve elevations of glycocalyx degradation products. Structural assessment using imaging methods show thinning of the endothelial glycocalyx layer in cardiovascular conditions and during major surgery, but thinning does not always correlate with the plasma concentrations of glycocalyx products. The few structural assessments performed do not currently support that capillary permeability is increased when the plasma levels of glycocalyx fragments in plasma are increased.

Conclusions: Shedding of glycocalyx components is a ubiquitous process that occurs during both acute and chronic inflammation with no sensitivity or specificity for a specific disease or condition.

1 | INTRODUCTION

The glycocalyx is a 0.2-5 µm thick layer of glycosylated proteins that covers the luminal side of the endothelium throughout the cardiovascular

system (Figure 1). The layer is believed to have great functional importance to local vasodilatation, coagulation, and inflammation.^{1,2}

Degradation of the endothelial glycocalyx layer ("shedding") occurs in inflammatory states, during ischemia, and after vigorous

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. Acta Anaesthesiologica Scandinavica published by John Wiley & Sons Ltd on behalf of Acta Anaesthesiologica Scandinavica Foundation.

volume loading. Such shedding is claimed to quickly change the physiology of the endothelium by obstructing local adaptation of blood flow and increasing the capillary permeability for macromolecules, which promotes hypovolemia.³

Given these properties, there is no wonder that the glycocalyx has received considerable attention over the past decade as demonstrated by the significant increase in publications. The majority of studies on glycocalyx shedding have been performed in laboratory animals, mostly in rats. Here, shedding of the endothelial glycocalyx seems to result primarily from metalloproteinase (MMP) activity, as inhibition of MMPs reduces endothelial glycocalyx degradation in response to inflammatory stimulation.⁴⁻⁶

Increases in circulating heparan sulfates, hyaluronan, and syndecan-1 have been reported in rat models of sepsis^{7,8} and hemorrhagic shock.⁹⁻¹¹

In terms of human investigations, the most common clinical study design has been to identify situations where the glycocalyx is degraded by measuring glycocalyx breakdown products in the plasma of patients. This has been done for a wide variety of conditions with the hypothesis being that the integrity of the glycocalyx can be related to changes in plasma concentrations of glycocalyx shed products.

In the discussion, we review confounding issues related to measuring plasma components assumed to represent cell-surface constituents but that have not been correlated to induce meaningful structural implications of clinical relevance.

2 | METHODS

To obtain a comprehensive dataset on all of the publications evaluating the human endothelial glycocalyx, a search of PubMed using the term "glycocalyx" yielded 3454 publications, returned from the earliest date in the database record, October 1958 to August 2020. The presentation adhered to the PRISMA Statement, whenever applicable.

Figure 2 represents a summary of our methodology. The first search was automated using the search terms "glycocalyx," then all

Editorial Comment


Glycocalyx shedding during inflammatory and ischemic states, and as a response to volume loading, is an very active area of research in recent years in anesthesiology and critical care medicine. This systematic review of glycocalyx shedding provides a detailed update on the current state of knowledge on quantification of glycocalyx shedding as assessed by measurements from plasma and in vivo structural measurements. Discussion is included regarding uncertainties when interpreting plasma measurements of breakdown products from glycocalyx shedding.

abstracts were manually reviewed by one person (VP) to ensure the Methods indicated that the study was: (a) human data, (b) recorded a glycocalyx component: syndecan-1 (sdc-1), syndecan-4 (sdc-4), glypican (Gpc), heparan sulfate (HS), chondroitin sulfate (CS) or hyaluronan (HA), and (c) recorded the source (plasma, CSF, urine). Every abstract published that included the word "glycocalyx" was reviewed in detail. The methods sections were reviewed for the type of study that was performed. Review articles and manuscripts without an Abstract in English or else did not evaluate the "human" glycocalyx, "glycocalyx integrity," and/or "glycocalyx damage" were excluded.

Structural assessments of the glycocalyx were categorized as sidestream dark field (SDF) imaging, orthogonal polarization spectral (OPS) imaging, incidental dark field (IDF) imaging, bright-field imaging and erythrocyte sodium sensitivity studies. A brief explanation of each method is presented in the Appendix. SDF, OPS, and IDF are quite similar methods but with incremental refinement to improve image clarity and resolution.

Abstracts that described structural assessment of the glycocalyx using fluorescent dextrans to determine void volumes of the glycocalyx were not included in this review due to concerns raised about this methodology.¹²

FIGURE 1 Schematic drawing of the molecular ultrastructure of the endothelial glycocalyx layer.² Used, by permission, from Bertram A, Stahl K, Hegemann J, Haller H. The glycocalyx layer. In: Hahn RG, ed. Clinical Fluid Therapy in the Perioperative Setting, 2nd Ed. Cambridge: Cambridge University Press [Colour figure can be viewed at wileyonlinelibrary.com]






FIGURE 2 Flow-chart showing the search strategy for the systematic review [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/ans.13797)]

We then categorized glycocalyx evaluation in common disease processes and organ systems studied by the authors. In doing so, we categorized common disease and syndromes as: critical illness; endocrine disease; pregnancy; surgery; healthy experimental patients; cardiovascular disease; renal disease; infectious disease; miscellaneous. From there, we identified subcategories in each category based on the models from the reviewed publications. These subcategories are given in the Tables that we chose to present our results.

After placing the reviewed publications into one of the major categories and subcategories, the publications were divided into three groups based on whether the publication investigated breakdown products, performed a structural assessment, or performed both.

The publications are listed in four tables using the following methods. We first assigned reference numbers in an ascending fashion to the models that performed an investigation of breakdown products as there were more publications that investigated this. These were assigned to Tables 1-3. We continued our reference list numerically to include the studies that performed a structural assessment. These were assigned to Table 4.

If a study performed both analysis of breakdown products and structural assessment, it was given the reference number that was labeled in Tables 1-3 and was also placed in Table 4 with the same numerical reference. We indicate an increase (+), decrease (-), or no change (0) in the glycocalyx breakdown product(s) depending on the individual study findings. Methods showing glycocalyx thickness were references as showing decrease (-), increase (+), or no change (0).

3 | RESULTS

After reviewing the 3,454 articles, secondary filters narrowed that list to 228 publications that met inclusion criteria of human studies.

Of these studies, 97% measured sdc-1, HS, and/or HA. The remaining studies assessed Gpc, CS, or synd-4.

3.1 | Commonly studied glycocalyx biomarkers

Sepsis and trauma are the most frequently studied conditions and comprise about 40 studies. They usually report 3-4-fold increased levels of glycocalyx degradation products. Exceptions are quite rare (Table 1, top). One study on encephalitis showed elevated concentrations of sdc-1 in the cerebrospinal fluid.⁶² By contrast, studies of diabetes more rarely report increased concentrations of glycocalyx degradation products. Among the few studies of pregnancy, an elevation is common when complications arise, such as preeclampsia or the HELLP syndrome (Table 1, bottom).

Surgery shows a variable picture. Major surgery that involves moments of ischemia and marked inflammation, such as cardiac surgery and transplantations, are most likely to show elevations of glycocalyx degradation products. The same number of studies of low-risk surgery show elevations as those that show no change or a decrease, which is the same overall picture offered by studies of healthy humans not undergoing surgery (Table 2).

Fourteen studies have measured glycocalyx degradation substances in cardiovascular disease, which is few when considering that the endothelium is part of the vascular system (Table 3, top). Heart failure and myocardial infarction show variable results. Other conditions show consistent elevations but are covered by single reports only. Renal disease shows a mixed picture, while all seven studies of infectious disease report elevations, mostly of syndecan-1. A number of other diseases in the internal medical field, such as lupus and leukemia, show consistent increases of glycocalyx degradation products. However, these conditions are mostly covered by single reports (Table 3, bottom).

TABLE 1 Changes in glyocalyx biomarker in critical illness, endocrine disease, and during pregnancy

Condition	Syndecan-1	Heparan sulfate	Hyaluronan	Source	Reference
Critical illness					
Sepsis	+			P	13-27
	+	+		P	28
	+		+	P	29,30
	-		-	P	31
		+	+	P	32,33
Trauma	+	+	+	P	34,36
		+	+	U	35 ^c
	+	+		P	37-52
	+	+	+	P	53
Traumatic Brain Injury	+			P	54 ^c
Stroke	+			P	55-58
Meningitis/ Encephalitis	+			(P, CSF)	59
	0			P	60
	0	+	0	P	61 ^{a,b,c}
Non-septic ICU/CCU	+			P	62
	+			CSF	62
	+			P	60,63
Endocrine disease	+	+		P	64,65
	-			P	66 ^a
Gestational diabetes	0	0	0	P	67 ^c
Type 1 diabetes		x		P	68,69
Type 2 diabetes	0			P	70
		x		P	71
Pregnancy					
Preeclampsia	-			P	72,73
	+			P	74,75
			+	P	76
	-		+	P	77
	0	+	+	P	78
C section	+	+		P	79
HEELP syndrome	+	+	+	P	80

Note: (P) plasma, (U) urine, (+) increase, (-) decrease, (0) no change

^aMeasured syndecan-4; ^bMeasured syndecan-2 and -3; ^cMeasured chondroitin sulfate

3.2 | Rarely studied glyocalyx biomarkers

Two studies included data on plasma Gpc; one study²¹ measured plasma from septic vs control patients and the second study¹⁶⁴ examined plasma glyocalyx markers (glypican and sdc-1) in relationship to forearm arterio-venous fistula failure. In sepsis, Gpc was increased along with C-reactive protein, lactate, pro-calcitonin, sdc-1, and heparin-binding protein. In the second study, there was no

relationship between plasma Gpc-1 or sdc-1 vs fistula failure; however, HA was positively correlated with failure of the arterio-venous fistula.

The five studies that reported on plasma concentrations of CS found increasing plasma concentrations during sepsis and acute respiratory failure,^{33,36} trauma,⁵⁴ ischemic stroke,⁶¹ and gestational diabetes.⁶⁷

The results for sdc-4 in plasma are more variable. One study found no change in sdc-4 during sepsis.³⁴ In a cohort of ICU patients

TABLE 2 Changes in glyocalyx biomarkers in surgical patients and healthy humans

Condition	Syndecan-1	Heparan Sulfate	Hyaluronan	Source	Reference
Surgery					
Cardiac surgery	+			P	81-88
	-			P	89-91
	0	0		P	92
	+	+		U	93
	+	+		P	94-97
	0	-	+	P	98
	+		+	P	99
	-		-	P	100
	+	0		P	101
	-			P	102
Vascular surgery	+	+		P	103,104
Abdominal surgery	+			P	105-107
	+	+		P	108
	0				109,110
	0	+	0	P	111
	0		0	P	112
	0	0	0	P	113,114
	-	-	+	U	114
	+			P	115-117
	+	+		P	118
	+		+	P	119
Thoracic surgery	+	-			120
	+			P	121
	+	0		P	122
	+	+		P	123
Elective low-risk surgery	-			P	124
	+		+	P	125
	+	0	+	P	126
	+	0		U	126
	+	0		P	127
	0	+	-	P	128
	0	-	-	U	128
Healthy humans ^b					
	+			P	129-134
	0				135
			+	P	136,137
		+	+	P	138
			0	P	139,140
	0	0	-	P	141
	+	+	0	P	142

Note: (P) plasma, (U) urine, (+) increase, (-) decrease, (0) no change

^bsubjected to exercise, simulated altitude, hypoxia etc

composed of gastrointestinal bleed, trauma, sepsis and cardiac arrest, sdc-4 was not different between patients groups or different from controls.²⁴ Sdc-4 decreased in ICU patients receiving lipid

emulsion infusion,⁶⁶ while an increase did occur in myocardial infarction.¹⁵¹ These limited data preclude any meaningful summary from being made.

TABLE 3 Changes in glyocalyx biomarkers in cardiovascular, infectious, and renal diseases

Condition	Syndecan-1	Heparan sulfate	Hyaluronan	Source	Reference
Cardiovascular					
Cardiomyopathy	+			P	143
Heart failure	+			P	144,145
	-	+		P	146
	0		+	P	147
Myocardial infarction	+			P	148-150
	0			P	151 ^a ,152
	+	+	+	P	
Coronary artery disease	+			P	154
	-			P	155
			+	P	156,157
Pulmonary embolism	+			P	158
Cardiac arrest	+		+	P	153,159
Renal					
Hemodialysis	+			P	160,161
	-			P	162
	+		+	P	163
	0		+	P	164
	+		+	P	165,166
Nephrotic syndrome	+				167,168
Acute kidney injury	+				169
	+	+	+		170
	-	-	-	U	170
Infectious					
	+			P	171-175
		+	+	P	176,177
	+	0	+	P	178 ^b
		+		U	179,180
Miscellaneous					
Leukemia	+			P	181
Lupus	+			P	182
Kawasaki's disease	+			P	183
	+		+	P	184
Vascular dementia			+	CSF	185
Transfusion-dependent	+			P	186-188
	+	0	0	P	189
Cirrhosis		+		P	190

Note: (P) plasma, (U) urine, (CSF) cerebrospinal fluid

(+) increase, (-) decrease, (0) no change

^asyndecan-4 increased; ^bchondroitin sulfate increased

3.3 | Structural measurements

The most widely used method for structural studies of the endothelial glyocalyx layer is to assess its thickness in the sublingual area by side Stream Dark Field (SDF) imaging. The assumption is that shedding of the glyocalyx layer both elevates the plasma concentration

of glyocalyx degradation products and causes a thinning or complete absence of this layer on the SDF image.

When sepsis has been studied by this approach, some studies have found a thinning of the glyocalyx layer while others have found no change (Table 4, top). There are two studies in stroke, and both also show a thinning.

TABLE 4 Structural measurements of glycocalyx thickness in various clinical settings. References with a lower number than 191 also report biomarkers, see Tables 1-3 for details

Condition	SDF	OPS	IDF	Other	Vascular bed	Ref.	Biomarker
Critical illness							
Sepsis	0				SL	191,192	
	-				SL	193-196	
				(-) Bright-field	SL	197	
Trauma			-		SL	38	sdC-1 +
Stroke	-				SL	198,199	
Non-septic ICU			-		SL, conjunctiva	63	sdC-1 +
	0					66	sdC-1 +
Endocrine disease							
Type 2 diabetes	+				SL and retinal	71	sdC-1 -
	+				SL	200	
	-				SL	202,203	
Type 1 diabetes		-			SL	68	sdC-1- HS- HA -
	-				SL	204,205	
Pregnancy							
Preeclampsia	-				SL	78	sdC-1- HS + HA +
Surgery							
Cardiac	-				SL	96,206,207	sdC-1 + HS +
	-				Skin	208	
Transplant surgery	-				Peritubular	118	sdC-1 + HA +
Elective low risk surgery	-				SL	209,210	
Neurosurgery	-					211	
				(-) Bright-field microscopy	SL	212	
Healthy humans ^a		-			SL	213	HS - HA -
				(-) Bright-field microscopy	Labial	214	
	+				SL	215-217	
	0				SL	218	HA +
	-				SL	219	
Cardiovascular							
Heart failure	-				SL	220	
	0				SL	221	
Hypertension	-				SL	222-224	
Coronary artery disease	-				SL	225,226	
	0				SL	227,228	
Pulmonary Artery Hypertension	0				SL	229	
Miscellaneous							
Hemodialysis				(+) sodium sensitivity	N/A	230	
				(-) sodium sensitivity	N/A	231	
	0				SL	232	sdC-1
Transfusion dependent	-				SL	189	sdC-1 + HA - HS -

(Continues)

TABLE 4 (Continued)

Condition	SDF	OPS	IDF	Other	Vascular bed	Ref.	Biomarker
Infection	-				Buccal	177	HS + HA +
	0				SL	233	
Antiphospholipid syndrome	-				SL	234	sdc-1+

Note: (SL) sublingual, (+) thickening, (-) reduction, (0) unchanged thickness

^asubjected to exercise, simulated altitude, hypoxia etc

Trauma and a stroke have been studied with Incidental Dark Field (IDF) imaging, and both show a thinning of the glycocalyx layer. Most studies of diabetes and all studies in major surgery also show thinning (Table 4, middle). The thickness of the glycocalyx in diabetes has also been studied with a third method, Orthogonal Polarization Spectral (OPS) imaging.

The results in healthy humans are quite mixed, while seven of the eight studies of cardiovascular conditions show thinning of the glycocalyx layer (Table 4, bottom).

Assessment of the glycocalyx thickness was made along with biomarker measurements in 15 studies of which 9 showed a correlation (60%). These studies can be found in Table 4.

4 | DISCUSSION

Quantification of glycocalyx shedding has become a common and expedient method to assess the integrity of the endothelial glycocalyx in human disease models. The number of publications increase at a high rate and involve many areas of human medicine. Sdc-1, HS, and HA are most commonly measured in humans, while Gpc, CS, and sdc-4 have been more rarely assessed. The many observational studies performed show that shedding of endothelial surface proteins is a common and ubiquitous occurrence during both acute and chronic inflammation. Less frequently, the thickness of the glycocalyx has been assessed by specialized imaging methods.

For anesthetists, the importance of the glycocalyx has been perceived to be functional and prognostic. The inflammation induced by trauma and major surgery triple the plasma concentrations of several molecules present in the glycocalyx, which has been attributed to endothelial injury. More pronounced elevations, averaging 15 times of baseline, has been reported after surgeries associated with ischemia, such as cardiac surgery.¹⁰⁵

The common pattern is that several glycocalyx biomarkers become elevated, but in some studies only sdc-1 was increased and not HS.^{101,122,126,178,189} This is surprising since syndecan-1 carries predominantly HS on its core protein and both markers would be expected to be correlated in plasma concentrations. However, this finding might be due to analytical error since syndecan-1 is stable in saved samples over time while heparan sulfate is more fragile, in particular if not consistently stored at -70°C until analyzed. Conversely, two studies reported a decrease in sdc-1 but an increase in HS or HA.^{77,146}

Elevated plasma concentrations of glycocalyx degradation products are clearly related to poor prognosis in severe disease, such as trauma^{44,47} and sepsis.^{29,105,196,235} However, our review still shows a paucity of data that links changes in plasma concentrations with quantitative physiological or pathological processes in humans. The sensitivity or specificity for examining glycocalyx fragments in plasma has not been demonstrated. A few studies show reductions in glycocalyx thickness but none of these studies make direct measurements of the functional consequences of such changes. Based on the Revised Starling Principle, glycocalyx shedding should increase the capillary permeability of macromolecules such as albumin, which reduces the plasma volume. We have found no studies in humans supporting that capillary permeability is increased when the plasma levels of glycocalyx fragments are increased. In fact, there is evidence that demonstrates no change in vascular permeability.^{65,109,236}

These uncertainties are unfortunate because clinical recommendations are frequently based on assumed connections of pathophysiological events involving the glycocalyx. For example, warnings have been given during the past decade that acute hypervolemia causes glycocalyx shedding and implicate that colloid fluids quickly lose most of their volume enhancing effect.²³⁷⁻²⁴⁰ Recent studies show no shedding from hypervolemia during surgery. These include cholecystitis, appendectomy,¹⁰⁹ hysterectomy,¹¹² and lengthy abdominal surgery.¹²⁸ Elevation of the sdc-1 and HA levels has been reported after correction for plasma dilution,¹²⁶ but the validity of such corrections is unproven due to a lack of pharmacokinetic characteristics for these substances.

Animal studies show that the glycocalyx layer might become degraded within 10 minutes, while restoration requires up to one week provided the shedding stimulus is removed.²⁴¹ In the rat lung restoration even occurs within 24 hours, but the recovery is dependent on the expression of the fibroblast growth factor receptor which is inhibited by sepsis.²⁴² These rates cannot be uncritically extrapolated to humans due to interspecies differences in metabolic rate and substrate turnover time.²⁴³

The metabolism of these compounds is very complicated, and with much of the elimination normally taking place in the liver. The kidneys are usually not considered to be of importance, but urinary concentrations actually do not deviate much from the plasma concentrations. A recent report highlighted the kidney as a source of elimination by measuring sdc-1, HS, and HA over 5 hours in healthy volunteers and post-surgical patients.¹⁷⁰ The renal clearance of

sdc-1 suggested that the entire plasma pool of free sdc-1 would be completely excreted within 15 hours and that a 6-fold variability in plasma concentration could be explained by acute changes in renal function and not due to increased shedding.

Glycocalyx constituents are assumed to stem from the endothelium despite being widely expressed in the body. Protein expression of sdc-1 is abundant in the liver, digestive tract, kidney, urinary bladder, and bone marrow, but hardly at all in muscle, adipose tissue, tongue, and skin. Protein expression for HS is found in muscle and occurs in the cytoplasm of many cell types, as well as in the interstitial matrix.²⁴⁴ It is often assumed that measured plasma constituents are uniformly derived from the luminal side of the endothelium throughout the vascular tree but, in fact, we do not know from where in the body they originate. This fact constitutes a bias in the present review, as authors being aware of the widespread distribution of glycosaminoglycans in the body may not always use manuscript titles and search terms that refer to the glycocalyx.²³⁵

We believe that direct measurement of perfused boundary region (PBR) in the microcirculation is, at present, the best method to assess degradation of the glycocalyx. The sublingual vessels have been used as an easily accessible vascular bed to determine the PBR, which is used a surrogate for glycocalyx thickness. In one study, sdc-1 was elevated but there were no changes in the PBR and glycocalyx dimensions obtained through structural imaging; Is the vascular endothelium the source, and do all vascular beds and endothelium contribute equally to shedding? Such questions are relevant because there are a number of human sepsis studies where sublingual PBR has no correlation with a variety of microvascular hemodynamic parameters.^{189,195} Specifically, PBR was not correlated with sublingual microvascular parameters including perfused vessel density, proportion of perfused vessels or microvascular flow index in resuscitated sepsis.¹⁹⁶ Likewise, PBR did not correlate with microvascular parameters in normodynamic vs hyperdynamic shock and there was limited correlation between the sublingual circulation and gut microcirculation in sepsis.^{189,245} Finally, this lack of correlation was reported following transfusion and use of activated protein C in septic patients.^{23,144} In summary, the current data show only a weak correlation between glycocalyx thickness and with structural microcirculatory indices during sepsis.

There is widespread acceptance of the glycocalyx as a permeability barrier. The original publication that gave rise to the glycocalyx' role as a permeability barrier came from Adamson who measured the hydraulic conductivity (L_p) of frog mesenteric microvessels before and after pronase treatment.²⁴⁶ Pronase is a broad-spectrum protease that was presumed to significantly degrade the glycocalyx off the surface of the mesenteric vessels. Adamson reported that pronase digestion of glycocalyx increased L_p by 2.5-fold and calculations estimated that glycocalyx accounted for 60% of hydraulic resistance to water flow across the capillary wall. However, increased capillary leakage of fluid or albumin resulting from increased plasma levels of sdc-1 and HS in humans has not been possible to demonstrate in volunteers and in surgical patients.^{65,109,110,236} Similarly, Ince *et al* and colleagues found no change in vascular barrier to fluid distribution

despite increases in plasma sdc-1 and HA in hemorrhaged rats.²⁴⁷ They also demonstrated that normovolemic hemodilution-induced glycocalyx shedding in rats does not alter vascular permeability to dextrans, albumin or plasma.²⁴⁸

In chronic diseases, compensatory responses may offset changes in shedding or expression. For example, in one study assessment of the glycocalyx by SDF imaging showed no significant difference in glycocalyx dimensions between patients with and without cardiovascular disease.²²⁷ However, two studies have reported elevations in sdc-1 and HA in patients with cardiovascular disease.^{159,184} Support for compensatory reaction(s) can be inferred from an *in vivo* study of the glycocalyx in sdc-1 knock-out (KO) mice where Savery *et al* used micro-particle velocimetry to determine the presence of a hydrodynamically significant cell surface layer.²⁴⁹ They observed no difference in wild-type vs. sdc-1 KO mice and concluded that sdc-1 was not necessary for the existence of an endothelial glycocalyx. Their limited analysis, however, could not rule out altered expression of other glycocalyx components in the sdc-1 KO. Consider that in cultured endothelial cells, knock down the sdc-1 gene resulted in increased mRNA for Gpc-1, sdc-4, sdc-4, and for a series of enzymes involved in heparan sulfate biosynthesis.²⁵⁰

Langford and colleagues examined the effect of mutated glucosaminoglycan attachment sites on the core protein of syndecan-1 on invasive cell function.²⁵¹ To their surprise, cells that expressed a modified syndecan with reduced glucosaminoglycan attachment sites on the core protein displayed no change in total cell surface heparan sulfate. This means the cells had compensatory responses to maintain a constant amount of HS on their surface to preserve function. If similar responses occur *in vivo* it is not surprising that sdc-1 KO mice have a normal glycocalyx. Evaluation of a single parameter then seems to be insufficient to make conclusion(s) about the composition, 3-dimensional structure, or functional consequences to the glycocalyx.

If we were to assume that all the data in Table 1 was free from publication bias and issues of renal clearance, the preponderance of data suggests that the glycocalyx is damaged during a wide variety of insults, as shedding increased in 96% of all published reports. If this percentage is valid, then what is the usefulness of a biomarker that increases ubiquitously across many disease states? Although uncertain, we suggest that an increase of five times the baseline in acute disease or trauma might be accepted as evidence of release of glycocalyx degradation products from somewhere in the body. Smaller changes might be due to short-term fluctuations in metabolism and urinary excretion.

5 | CONCLUSION

The utility of measuring glycocalyx breakdown products in the plasma as biomarkers with a predictive value for determining a specific disease or the progression of a disease is unproven. Shedding of glycocalyx components is a ubiquitous process that occurs during both acute and chronic inflammation with no sensitivity or specificity

for a specific disease or condition. Uncertainties related to proteoglycan expression levels, turnover rate, shedding, renal clearance and lack of correction for hemodilution cast doubts on many of the reported alterations in glycocalyx breakdown products measured in plasma. There is only a moderately good correlation between plasma concentration and the structural assessment of glycocalyx thickness (60% agreement), which further questions the utility of measuring plasma glycocalyx components as a surrogate for structural and functional alterations. Finally, compensatory expression of glycocalyx constituents may offset the loss of specific components in order to maintain structural integrity.

CONFLICT OF INTEREST

None.

ORCID

Robert G. Hahn  <https://orcid.org/0000-0002-1528-3803>

REFERENCES

- Pries AR, Secomb TW, Gaehtgens P. The endothelial surface layer. *Pflügers Arch.* 2000;440:653-666.
- Bertram A, Stahl K, Hegemann J, Haller H. The glycocalyx layer. In: Hahn RG, ed. *Clinical Fluid Therapy in the Perioperative Setting*, 2nd edn. Cambridge: Cambridge University Press; 2016:73-81.
- Alphonsus CS, Rodseth RN. The endothelial glycocalyx: a review of the vascular barrier. *Anaesthesia.* 2014;69:777-784.
- Cui N, Wang H, Long Y, Su L, Liu D. Dexamethasone suppressed LPS-induced matrix metalloproteinase and its effect on endothelial glycocalyx shedding. *Mediators Inflamm.* 2015;2015:1-8.
- Lipowsky HH, Lescanic A, Sah R. Role of matrix metalloproteinases in the kinetics of leucocyte-endothelial adhesion in post-capillary venules. *Biorheol.* 2015;52:433-445.
- Lipowsky HH, Sah R, Lescanic A. Relative roles of doxycycline and cation chelation in endothelial glycan shedding and adhesion of leukocytes. *Am J Physiol Heart Circ Physiol.* 2011;300:H415-H422.
- Lipowsky HH, Gao L, Lescanic A. Shedding of the endothelial glycocalyx in arterioles, capillaries, and venules and its effect on capillary hemodynamics during inflammation. *Am J Physiol Heart Circ Physiol.* 2011;301:H2235-H2245.
- Huang X, Han S, Liu X, et al. Both UFH and NAH alleviate shedding of endothelial glycocalyx and coagulopathy in LPS-induced sepsis. *Exp Ther Med.* 2020;19:913-922.
- Torres Filho IP, Torres LN, Salgado C, Dubick MA. Plasma syndecan-1 and heparan sulfate correlate with microvascular glycocalyx degradation in hemorrhaged rats after different resuscitation fluids. *Am J Physiol Heart Circ Physiol.* 2016;310:H1468-H1478.
- Wu F, Peng Z, Park PW, Kozar RA. Loss of syndecan-1 abrogates the pulmonary protective phenotype induced by plasma after hemorrhagic shock. *Shock.* 2017;48:340-345.
- Kozar RA, Peng Z, Zhang R, et al. Plasma restoration of endothelial glycocalyx in a rodent model of hemorrhagic shock. *Anesth Analg.* 2011;112:1289-1295.
- Michel CC, Curry FR. Glycocalyx volume: a critical review of tracer dilution methods for its measurement. *Microcirculation.* 2009;16(3):213-219. <https://doi.org/10.1080/10739680802527404>
- Ikeda M, Matsumoto H, Ogura H, et al. Circulating syndecan-1 predicts the development of disseminated intravascular coagulation in patients with sepsis. *J Crit Care.* 2018;43:48-153.
- Wu X, Hu Z, Yuan H, Che L, Li Y, Zhao C. Fluid resuscitation and markers of glycocalyx degradation in severe sepsis. *Open Med (Wars).* 2017;12(1):409-416. <https://doi.org/10.1515/med-2017-0059>
- Murphy LS, Wickersham N, McNeil JB, et al. Endothelial glycocalyx degradation is more severe in patients with non-pulmonary sepsis compared to pulmonary sepsis and associates with risk of ARDS and other organ dysfunction. *Ann Intensive Care.* 2017;7:102.
- Puskarich MA, Cornelius DC, Tharp J, Nandi U, Jones AE. Plasma syndecan-1 levels identify a cohort of patients with severe sepsis at high risk for intubation after large-volume intravenous fluid resuscitation. *J Crit Care.* 2016;36:125-129.
- Ostrowski SR, Gaini S, Pedersen C, Johansson PI. Sympathoadrenal activation and endothelial damage in patients with varying degrees of acute infectious disease: an observational study. *J Crit Care.* 2015;30:90-96.
- Johansson PI, Haase N, Perner A, Ostrowski SR. Association between sympathoadrenal activation, fibrinolysis, and endothelial damage in septic patients: a prospective study. *J Crit Care.* 2014;29:327-333.
- Ostrowski SR, Berg RM, Windelov NA, et al. Coagulopathy, catecholamines, and biomarkers of endothelial damage in experimental human endotoxemia and in patients with severe sepsis: a prospective study. *J Crit Care.* 2013;28:586-596.
- Sallissalmi M, Tenhunen J, Yang R, Oksala N, Pettila V. Vascular adhesion protein-1 and syndecan-1 in septic shock. *Acta Anaesthesiol Scand.* 2012;56:316-322.
- Fisher J, Linder A, Bentzer P. Elevated plasma glypicans are associated with organ failure in patients with infection. *Intensive Care Med Exp.* 2019;7:2.
- Wei S, Gonzalez Rodriguez E, Chang R, et al. Elevated syndecan-1 after trauma and risk of sepsis: a secondary analysis of patients from the pragmatic, randomized optimal platelet and plasma ratios (PROPPR) trial. *J Am Coll Surg.* 2018;227:587-595.
- Johansson PI, Henriksen HH, Stensballe J, et al. Traumatic endotheliopathy: a prospective observational study of 424 severely injured patients. *Ann Surg.* 2017;265:597-603.
- Nelson A, Johansson J, Tyden J, Bodelsson M. Circulating syndecans during critical illness. *APMIS.* 2017;125:468-475.
- de Oliveira Neves FM, Araujo CB, de Freitas DF, et al. Fibroblast growth factor 23, endothelium biomarkers and acute kidney injury in critically-ill patients. *J Transl Med.* 2019;17:121.
- Welling H, Henriksen HH, Gonzalez-Rodriguez ER, et al. Endothelial glycocalyx shedding in patients with burns. *Burns.* 2020;2020(46):386-393.
- Nelson A, Berkestedt I, Schmidtchen A, Ljunggren L, Bodelsson M. Increased levels of glycosaminoglycans during septic shock: relation to mortality and the antibacterial actions of plasma. *Shock.* 2008;30:623-627.
- Hippensteel JA, Uchimido R, Tyler PD, et al. Intravenous fluid resuscitation is associated with septic endothelial glycocalyx degradation. *Crit Care.* 2019;23:259.
- Anand D, Ray S, Srivastava LM, Bhargava S. Evolution of serum hyaluronan and syndecan levels in prognosis of sepsis patients. *Clin Biochem.* 2016;49:768-776.
- Smart L, Bosio E, Macdonald SPJ, et al. Glycocalyx biomarker syndecan-1 is a stronger predictor of respiratory failure in patients with sepsis due to pneumonia, compared to endocan. *J Crit Care.* 2018;47:93-98.
- Smart L, Macdonald SPJ, Bosio E, Fatovich D, Neil C, Arendts G. Bolus therapy with 3% hypertonic saline or 0.9% saline in emergency department patients with suspected sepsis: a pilot randomized controlled trial. *J Crit Care.* 2018;52:33-39.
- Li M, Hao D, Wang T, et al. Variation and clinical value of endothelial glycocalyx in the patients with septic shock. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue.* 2016;28:699-703.
- Schmidt EP, Li G, Li L, et al. The circulating glycosaminoglycan signature of respiratory failure in critically ill adults. *J Biol Chem.* 2014;289:8194-8202.

34. Smart L, Macdonald SPJ, Burrows S, Bosio E, Arendts G, Fatovich DM. Endothelial glycocalyx biomarkers increase in patients with infection during emergency department treatment. *J Crit Care*. 2017;42:304-309.
35. Schmidt EP, Overdier KH, Sun X, et al. Urinary glycosaminoglycans predict outcomes in septic shock and acute respiratory distress syndrome. *Am J Respir Crit Care Med*. 2016;194:439-449.
36. Huang X, Hu H, Sun T, et al. Plasma endothelial glycocalyx components as a potential biomarker for predicting the development of disseminated intravascular coagulation in patients with sepsis. *J Intensive Care Med*. 2020;885066620949131: <https://doi.org/10.1177/0885066620949131>
37. Russell RT, Christiaans SC, Nice TR, et al. Histone-complexed DNA fragments levels are associated with coagulopathy, endothelial cell damage, and increased mortality after severe pediatric trauma. *Shock*. 2018;49:44-52.
38. Naumann DN, Hazeldine J, Midwinter MJ, Hutchings SD, Harrison P. Poor microcirculatory flow dynamics are associated with endothelial cell damage and glycocalyx shedding after traumatic hemorrhagic shock. *J Trauma Acute Care Surg*. 2018;84:81-88.
39. Rodriguez EG, Cardenas JC, Lopez E, et al. Early identification of the patient with endotheliopathy of trauma by arrival serum albumin. *Shock*. 2017;50:31-37.
40. Ostrowski SR, Henriksen HH, Stensballe J, et al. Sympathoadrenal activation and endotheliopathy are drivers of hypocoagulability and hyperfibrinolysis in trauma: a prospective observational study of 404 severely injured patients. *J Trauma Acute Care Surg*. 2017;82:293-301.
41. Gonzalez Rodriguez E, Ostrowski SR, Cardenas JC, et al. Syndecan-1: a quantitative marker for the endotheliopathy of trauma. *J Am Coll Surg*. 2017;225:419-427.
42. Ostrowski SR, Sørensen AM, Windeløv NA, et al. High levels of soluble VEGF receptor 1 early after trauma are associated with shock, sympathoadrenal activation, glycocalyx degradation and inflammation in severely injured patients: a prospective study. *Scand J Trauma Resusc Emerg Med*. 2012;20:27.
43. Ostrowski SR, Johansson PI. Endothelial glycocalyx degradation induces endogenous heparinization in patients with severe injury and early traumatic coagulopathy. *J Trauma Acute Care Surg*. 2012;73:60-66.
44. Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR. High circulating adrenaline levels at admission predict increased mortality after trauma. *J Trauma Acute Care Surg*. 2012;72:428-436.
45. Johansson PI, Sorensen A, Perner A, et al. High sCD40L levels early after trauma are associated with enhanced shock, sympathoadrenal activation, tissue and endothelial damage, coagulopathy and mortality. *J Thromb Haemost*. 2012;10:207-216.
46. Johansson PI, Sorensen AM, Perner A, et al. Elderly trauma patients have high circulating noradrenaline levels but attenuated release of adrenaline, platelets, and leukocytes in response to increasing injury severity. *Crit Care Med*. 2012;40:1844-1850.
47. Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR. A high admission syndecan-1 level, a marker of endothelial glycocalyx degradation, is associated with inflammation, protein C depletion, fibrinolysis, and increased mortality in trauma patients. *Ann Surg*. 2011;254:194-200.
48. Johansson PI, Sorensen AM, Perner A, et al. Disseminated intravascular coagulation or acute coagulopathy of trauma shock early after trauma? An observational study. *Crit Care*. 2011;15:R272.
49. Wade CE, Matijevic N, Wang YW, et al. Absences of endothelial microvesicle changes in the presence of the endotheliopathy of trauma. *Shock*. 2019;51:180-184.
50. Russell RT, McDaniel JK, Cao W, et al. Low plasma ADAMTS13 activity is associated with coagulopathy, endothelial cell damage and mortality after severe paediatric trauma. *Thromb Haemost*. 2018;118:676-687.
51. Richter RP, Russell RT, Hu PJ, et al. Plasma angiopoietin-2/-1 ratio is elevated and angiopoietin-2 levels correlate with plasma syndecan-1 following pediatric trauma. *Shock*. 2019;52:340-346.
52. Henriksen HH, McGarrity S, Sigurðardóttir RS, et al. Metabolic systems analysis of shock-induced endotheliopathy (SHINE) in trauma: a new research paradigm. *Ann Surg*. 2019;272:1140-1148.
53. Halbgebauer R, Braun CK, Denk S, et al. Hemorrhagic shock drives glycocalyx, barrier and organ dysfunction early after polytrauma. *J Crit Care*. 2018;44:229-237.
54. Rahbar E, Cardenas JC, Baimukanova G, et al. Endothelial glycocalyx shedding and vascular permeability in severely injured trauma patients. *J Transl Med*. 2015;13:117.
55. Albert V, Subramanian A, Agrawal D, Pati HP, Gupta SD, Mukhopadhyay AK. Acute Traumatic endotheliopathy in isolated severe brain injury and its impact on clinical outcome. *Med Sci (Basel)*. 2018;6:5.
56. Di Battista AP, Rizoli SB, Lejnieks B, et al. Sympathoadrenal activation is associated with acute traumatic coagulopathy and endotheliopathy in isolated brain injury. *Shock*. 2016;46:96-103.
57. Genet GF, Johansson PI, Meyer MA, et al. Trauma-induced coagulopathy: standard coagulation tests, biomarkers of coagulopathy, and endothelial damage in patients with traumatic brain injury. *J Neurotrauma*. 2013;30:301-306.
58. Gonzalez Rodriguez E, Cardenas JC, Cox CS, et al. Traumatic brain injury is associated with increased syndecan-1 shedding in severely injured patients. *Scand J Trauma Resusc Emerg Med*. 2018;26(1):102. <https://doi.org/10.1186/s13049-018-0565-3>
59. Bell JD, Rhind SG, Di Battista AP, Macdonald RL, Baker AJ. Biomarkers of glycocalyx injury are associated with delayed cerebral ischemia following aneurysmal subarachnoid hemorrhage: a case series supporting a new hypothesis. *Neurocrit Care*. 2017;26(3):339-347. <https://doi.org/10.1007/s12028-016-0357-4>
60. Gybel-Brask M, Rasmussen R, Stensballe J, Johansson PI, Ostrowski SR. Effect of delayed onset prostacyclin on markers of endothelial function and damage after subarachnoid hemorrhage. *Acta Neurochir (Wien)*. 2017;159(6):1073-1078. <https://doi.org/10.1007/s00701-017-3168-2>
61. DellaValle B, Hasseldam H, Johansen FF, Iversen HK, Rungby J, Hempel C. Multiple soluble components of the glycocalyx are increased in patient plasma after ischemic stroke. *Stroke*. 2019;50(10):2948-2951. <https://doi.org/10.1161/STROKEAHA.119.025953>
62. Zhu J, Li Y, Zheng D, et al. Elevated serum and cerebrospinal fluid CD138 in patients with anti-N-Methyl-D-Aspartate receptor encephalitis. *Front Mol Neurosci*. 2019;12:116. <https://doi.org/10.3389/fnmol.2019.00116>
63. Pranskunas A, Tamosiutis T, Balciuniene N, et al. Alterations of conjunctival glycocalyx and microcirculation in non-septic critically ill patients. *Microvasc Res*. 2018;118:44-48.
64. Osuka A, Kusuki H, Yoneda K, et al. Glycocalyx shedding is enhanced by age and correlates with increased fluid requirement in patients with major burns. *Shock*. 2018;50(1):60-65. <https://doi.org/10.1097/SHK.0000000000001028>
65. Hahn RG, Zdolsek M, Krizhanovskii C, Ntika S, Zdolsek J. Elevated plasma concentrations of syndecan-1 do not correlate with increased capillary leakage of 20% albumin. *Anesth Analg*. 2020;132:856-865. <https://doi.org/10.1213/ANE.00000000000051315>
66. Astapenko D, Turek Z, Dostal P, et al. Effect of short-term administration of lipid emulsion on endothelial glycocalyx integrity in ICU patients - a microvascular and biochemical pilot study. *Clin Hemorheol Microcirc*. 2019;73(2):329-339. <https://doi.org/10.3233/CH-190564>

67. Long DS, Hou W, Taylor RS, McCowan LM. Serum levels of endothelial glycocalyx constituents in women at 20 weeks' gestation who later develop gestational diabetes mellitus compared to matched controls: a pilot study. *BMJ Open*. 2016;6:e011244. <https://doi.org/10.1136/bmjopen-2016-011244>
68. Nieuwdorp M, Holleman F, de Groot E, et al. Perturbation of hyaluronan metabolism predisposes patients with type 1 diabetes mellitus to atherosclerosis. *Diabetologia*. 2007;50(6):1288-1293. <https://doi.org/10.1007/s00125-007-0666-4>
69. Nieuwdorp M, Mooij HL, Kroon J, et al. Endothelial glycocalyx damage coincides with microalbuminuria in type 1 diabetes. *Diabetes*. 2006;55(4):1127-1132. <https://doi.org/10.2337/diabetes.55.04.06.db05-1619>
70. Nassimzadeh M, Ashrafian H, Drury NE, et al. Reduced negative surface charge on arterial endothelium explains accelerated atherosclerosis in type 2 diabetic patients. *Diab Vasc Dis Res*. 2010;7(3):213-215. <https://doi.org/10.1177/1479164110376207>
71. Broekhuizen LN, Lemkes BA, Mooij HL, et al. Effect of sulodexide on endothelial glycocalyx and vascular permeability in patients with type 2 diabetes mellitus. *Diabetologia*. 2010;53(12):2646-2655. <https://doi.org/10.1007/s00125-010-1910-x>
72. Gandle RE, Althouse A, Jayabalan A, et al. Low soluble syndecan-1 precedes preeclampsia. *PLoS One*. 2016;11(6):e0157608. <https://doi.org/10.1371/journal.pone.0157608>
73. Bramham K, Villa PM, Joslin JR, et al. Predisposition to superimposed preeclampsia in women with chronic hypertension: endothelial, renal, cardiac, and placental factors in a prospective longitudinal cohort. *Hypertens Pregnancy*. 2020;39(3):326-335. <https://doi.org/10.1080/10641955.2020.1769643>
74. Kuessel L, Husslein H, Montanari E, et al. Dynamics of soluble syndecan-1 in maternal serum during and after pregnancies complicated by preeclampsia: a nested case control study. *Clin Chem Lab Med*. 2019;58(1):50-58. <https://doi.org/10.1515/cclm-2019-0686>
75. Greeley ET, Rochelson B, Krantz DA, et al. Evaluation of syndecan-1 as a novel biomarker for adverse pregnancy outcomes. *Reprod Sci*. 2020;27:355-363.
76. Wiles K, Bramham K, Seed PT, et al. Diagnostic indicators of superimposed preeclampsia in women with CKD. *Kidney Int Rep*. 2019;4(6):842-853. <https://doi.org/10.1016/j.ekir.2019.03.012>
77. Kornacki J, Wirstlein P, Wender-Ozegowska E. Levels of syndecan-1 and hyaluronan in early- and late-onset preeclampsia. *Pregnancy Hypertens*. 2019;18:108-111.
78. Weissgerber TL, Garcia-Valencia O, Milic NM, et al. Early onset preeclampsia is associated with glycocalyx degradation and reduced microvascular perfusion. *J Am Heart Assoc*. 2019;8:e010647. <https://doi.org/10.1161/JAHA.118.010647>
79. Powell MF, Mathru M, Brandon A, Patel R, Frolich MA. Assessment of endothelial glycocalyx disruption in term parturients receiving a fluid bolus before spinal anesthesia: a prospective observational study. *Int J Obstet Anesth*. 2014;23(4):330-334. <https://doi.org/10.1016/j.ijoa.2014.06.001>
80. Hofmann-Kiefer KF, Knabl J, Martinoff N, et al. Increased serum concentrations of circulating glycocalyx components in HELLP syndrome compared to healthy pregnancy: an observational study. *Reprod Sci*. 2013;20(3):318-325. <https://doi.org/10.1177/1933719112453508>
81. de Melo Bezerra Cavalcante CT, Castelo Branco KM, Pinto Júnior VC, et al. Syndecan-1 improves severe acute kidney injury prediction after pediatric cardiac surgery. *J Thorac Cardiovasc Surg*. 2016;152(1):178-186. <https://doi.org/10.1016/j.jtcvs.2016.03.079>
82. Ostrowski SR, Pedersen SH, Jensen JS, Mogelvang R, Johansson PI. Acute myocardial infarction is associated with endothelial glycocalyx and cell damage and a parallel increase in circulating catecholamines. *Crit Care*. 2013;17(1):R32. <https://doi.org/10.1186/cc12532>
83. Idorn L, Jensen AS, Juul K, et al. Thromboembolic complications in Fontan patients: population-based prevalence and exploration of the etiology. *Pediatr Cardiol*. 2013;34(2):262-272. <https://doi.org/10.1007/s00246-012-0431-4>
84. Svennevig K, Hoel TN, Thiara AS, et al. Syndecan-1 plasma levels during coronary artery bypass surgery with and without cardiopulmonary bypass. *Perfusion*. 2008;23(3):165-171. <https://doi.org/10.1177/0267659108098215>
85. Pesonen E, Passov A, Andersson S, et al. Glycocalyx degradation and inflammation in cardiac surgery. *J Cardiothorac Vasc Anesth*. 2019;33(2):341-345. <https://doi.org/10.1053/j.jvca.2018.04.007>
86. Dekker NAM, Veerhoek D, van Leeuwen ALI, Vonk ABA, van den Brom CE, Boer C. Microvascular alterations during cardiac surgery using a heparin or phosphorylcholine-coated circuit. *J Cardiothorac Vasc Anesth*. 2020;34:912-919.
87. Robich M, Ryzhov S, Kacer D, et al. Prolonged cardiopulmonary bypass is associated with endothelial glycocalyx degradation. *J Surg Res*. 2020;251:287-295.
88. Kim HB, Soh S, Kwak YL, Bae JC, Kang SH, Song JW. High Preoperative serum syndecan-1, a marker of endothelial glycocalyx degradation, and severe acute kidney injury after valvular heart surgery. *J Clin Med*. 2020;9:1803.
89. Pesonen E, Keski-Nisula J, Andersson S, Palo R, Salminen J, Suominen PK. High-dose methylprednisolone and endothelial glycocalyx in paediatric heart surgery. *Acta Anaesthesiol Scand*. 2016;60(10):1386-1394. <https://doi.org/10.1111/aas.12785>
90. Holmvang L, Ostrowski SR, Dridi NP, Johansson P. A single center, open, randomized study investigating the clinical safety and the endothelial modulating effects of a prostacyclin analog in combination with eptifibatide in patients having undergone primary percutaneous coronary intervention (PCI) for ST-segment elevation myocardial infarction. *Prostaglandins Other Lipid Mediat*. 2012;99(3-4):87-95. <https://doi.org/10.1016/j.prostaglandins.2012.08.002>
91. Kim TK, Nam K, Cho YJ, et al. Microvascular reactivity and endothelial glycocalyx degradation when administering hydroxyethyl starch or crystalloid during off-pump coronary artery bypass graft surgery: a randomised trial. *Anaesthesia*. 2017;72(2):204-213. <https://doi.org/10.1111/anae.13642>
92. Abou-Arab O, Kamel S, Beyls C, et al. Vasoplegia after cardiac surgery is associated with endothelial glycocalyx alterations. *J Cardiothorac Vasc Anesth*. 2019;34:900-905. <https://doi.org/10.1053/j.jvca.2019.09.004>
93. Ferrer NMB, de Melo Bezerra Cavalcante CT, Branco KMC, et al. Urinary Syndecan-1 and acute kidney injury after pediatric cardiac surgery. *Clin Chim Acta*. 2018;485:205-209. <https://doi.org/10.1016/j.cca.2018.06.033>
94. Bruegger D, Rehm M, Abicht J, et al. Shedding of the endothelial glycocalyx during cardiac surgery: on-pump versus off-pump coronary artery bypass graft surgery. *J Thorac Cardiovasc Surg*. 2009;138(6):1445-1447. <https://doi.org/10.1016/j.jtcvs.2008.07.063>
95. Brettner F, Chappell D, Nebelsiek T, et al. Preinterventional hydrocortisone sustains the endothelial glycocalyx in cardiac surgery. *Clin Hemorheol Microcirc*. 2019;71(1):59-70. <https://doi.org/10.3233/CH-180384>
96. Dekker NAM, Veerhoek D, Koning NJ, et al. Postoperative microcirculatory perfusion and endothelial glycocalyx shedding following cardiac surgery with cardiopulmonary bypass. *Anaesthesia*. 2019;74(5):609-618. <https://doi.org/10.1111/anae.14577>
97. He G, Gao Y, Feng L, et al. Correlation between wall shear stress and acute degradation of the endothelial glycocalyx during cardiopulmonary bypass. *J Cardiovasc Transl Res*. 2020;13(6):1024-1032. <https://doi.org/10.1007/s12265-020-10027-2>
98. Hohn A, Baumann A, Pietroschinsky E, et al. Haemoadsorption: effective in reducing circulating fragments of the endothelial

- glycocalyx during cardiopulmonary bypass in patients undergoing on-pump cardiac surgery? *Minerva Anesthesiol.* 2020;87:35-42. <https://doi.org/10.23736/S0375-9393.20.14525-5>
99. Bruegger D, Brettner F, Rossberg I, et al. Acute degradation of the endothelial glycocalyx in infants undergoing cardiac surgical procedures. *Ann Thorac Surg.* 2015;99(3):926-931. <https://doi.org/10.1016/j.athoracsur.2014.10.013>
 100. Mennander AA, Shalaby A, Oksala N, et al. Diazoxide may protect endothelial glycocalyx integrity during coronary artery bypass grafting. *Scand Cardiovasc J.* 2012;46(6):339-344. <https://doi.org/10.3109/14017431.2012.717303>
 101. Li X, Sun S, Wu G, Che X, Zhang J. Effect of Hydroxyethyl Starch Loading on Glycocalyx shedding and cerebral metabolism during surgery. *J Surg Res.* 2020;246:274-283. <https://doi.org/10.1016/j.jss.2019.09.030>
 102. Stensballe J, Ulrich AG, Nilsson JC, et al. Resuscitation of endotheliopathy and bleeding in thoracic aortic dissections: The VIPER-OCTA Randomized Clinical Pilot Trial. *Anesth Analg.* 2018;127(4):920-927. <https://doi.org/10.1213/ANE.00000000000003545>
 103. Martin L, Gombert A, Chen J, et al. The beta-d-endoglucuronidase heparanase is a danger molecule that drives systemic inflammation and correlates with clinical course after open and endovascular thoracoabdominal aortic aneurysm repair: lessons learnt from mice and men. *Front Immunol.* 2017;8:681. <https://doi.org/10.3389/fimmu.2017.00681>
 104. Rehm M, Bruegger D, Christ F, et al. Shedding of the endothelial glycocalyx in patients undergoing major vascular surgery with global and regional ischemia. *Circulation.* 2007;116:1896-1906.
 105. Holzmans MS, Winkler MS, Strunden MS, et al. Syndecan-1 as a biomarker for sepsis survival after major abdominal surgery. *Biomark Med.* 2018;12:119-127.
 106. Wang JW, Wu AS, Yue Y, Wu Y. Perioperative ulinastatin helps preserve endothelial glycocalyx layer in periampullary carcinoma patients undergoing traditional whipple procedure. *Clin Hemorheol Microcir.* 2020;75(2):135-142. <https://doi.org/10.3233/CH-190688>
 107. Pustetto M, Goldsztejn N, Touhri K, Engelman E, Ickx B, van Obbergh L. Intravenous lidocaine to prevent endothelial dysfunction after major abdominal surgery: a randomized controlled pilot trial. *BMC Anesthesiol.* 2020;20:155.
 108. Steppan J, Hofer S, Funke B, et al. Sepsis and major abdominal surgery lead to flaking of the endothelial glycocalyx. *J Surg Res.* 2011;165:136-141.
 109. Li Y, Yi S, Zhu Y, Hahn RG. Volume kinetics of Ringer's lactate solution in acute inflammatory disease. *Br J Anaesth.* 2018;121:574-580.
 110. Statkevicius S, Bonnevier J, Fisher J, et al. Albumin infusion rate and plasma volume expansion: a randomized clinical trial in postoperative patients after major surgery. *Crit Care.* 2019;23:191.
 111. Kammerer T, Hulde N, Speck E, et al. Effects of balanced hydroxyethyl starch 6% (130/0.4) and albumin 5% on clot formation and glycocalyx shedding: Subgroup analysis of a prospective randomized trial. *Thromb Res.* 2019;183:111-118.
 112. Nemme J, Krizhanovskii C, Ntikia S, Sabelnikovs O, Vanags I, Hahn RG. Hypervolaemia does not cause shedding of the endothelial glycocalyx layer during hysterectomy; a randomised clinical trial comparing sevoflurane and propofol anaesthesia. *Acta Anaesthesiol Scand.* 2020;64:538-545.
 113. Kaneko T, Tataru T, Hirose M. Effects of anaesthesia-induced hypotension and phenylephrine on plasma volume expansion by hydroxyethyl starch: a randomised controlled study. *Acta Anaesthesiol Scand.* 2020;64:620-627.
 114. Nemme J, Hahn RG, Krizhanovskii C, Ntika S, Sabelnikovs O, Vanags I. Minimal shedding of the glycocalyx layer during abdominal hysterectomy. *BMC Anesthesiol.* 2017;17:107.
 115. Schiefer J, Lebherz-Eichinger D, Erdoes G, et al. Alterations of endothelial glycocalyx during orthotopic liver transplantation in patients with end-stage liver disease. *Transplantation.* 2015;99:2118-2123.
 116. Adepu S, Rosman CWK, Dam W, et al. Incipient renal transplant dysfunction associates with tubular syndecan-1 expression and shedding. *Am J Physiol Renal Physiol.* 2015;309:F137-F145.
 117. Schiefer J, Faybik P, Koch S, et al. Glycocalyx damage within human liver grafts correlates with graft injury and postoperative graft function after orthotopic liver transplantation. *Transplantation.* 2020;104:72-78.
 118. Snoeijs MG, Vink H, Voesten N, et al. Acute ischemic injury to the renal microvasculature in human kidney transplantation. *Am J Physiol Renal Physiol.* 2010;299:F1134-F1140.
 119. Sladden TM, Yerkovich S, Grant M, et al. Endothelial glycocalyx shedding predicts donor organ acceptability and is associated with primary graft dysfunction in lung transplant recipients. *Transplantation.* 2019;103:1277-1285.
 120. Passov A, Schramko A, Makisalo H, et al. Graft glycocalyx degradation in human liver transplantation. *PLoS One.* 2019;14(8):e0221010. <https://doi.org/10.1371/journal.pone.0221010>
 121. Wang J, Wu A, Wu Y. Endothelial glycocalyx layer: a possible therapeutic target for acute lung injury during lung resection. *Biomed Res Int.* 2017;2017:1-8. <https://doi.org/10.1155/2017/5969657>
 122. Arthur A, McCall PJ, Jolly L, Kinsella J, Kirk A, Shelley BG. Endothelial glycocalyx layer shedding following lung resection. *Biomark Med.* 2016;10(10):1033-1038. <https://doi.org/10.2217/bmm-2016-0163>
 123. Kim HJ, Kim E, Baek SH, et al. Sevoflurane did not show better protective effect on endothelial glycocalyx layer compared to propofol during lung resection surgery with one lung ventilation. *J Thorac Dis.* 2018;10:1468-1475.
 124. Lindberg-Larsen V, Ostrowski SR, Lindberg-Larsen M, Rovsing ML, Johansson PI, Kehlet H. The effect of pre-operative methylprednisolone on early endothelial damage after total knee arthroplasty: a randomised, double-blind, placebo-controlled trial. *Anaesthesia.* 2017;72:1217-1224.
 125. Belavic M, Sotosek Tokmadzic V, Fistic E, et al. The effect of various doses of infusion solutions on the endothelial glycocalyx layer in laparoscopic cholecystectomy patients. *Minerva Anesthesiol.* 2018;84:1032-1043.
 126. Chappell D, Bruegger D, Potzel J, et al. Hypervolemia increases release of atrial natriuretic peptide and shedding of the endothelial glycocalyx. *Crit Care.* 2014;18:538.
 127. Maldonado F, Morales D, Gutiérrez R, Barahona M, Cerda O, Cáceres M. Effect of sevoflurane and propofol on tourniquet-induced endothelial damage: a pilot randomized controlled trial for knee-ligament surgery. *BMC Anesthesiol.* 2020;20:121.
 128. Hasselgren E, Zdosek M, Zdosek JH, et al. Long intravascular persistence of 20% albumin in postoperative patients. *Anesth Analg.* 2019;129:1232-1239.
 129. Johansson PI, Bergstrom A, Aachmann-Andersen NJ, et al. Effect of acute hypobaric hypoxia on the endothelial glycocalyx and digital reactive hyperemia in humans. *Front Physiol.* 2014;5:459.
 130. Bihari S, Wiersema UF, Perry R, et al. Efficacy and safety of 20% albumin fluid loading in healthy subjects: a comparison of four resuscitation fluids. *J Appl Physiol.* 2019;126:1646-1660.
 131. Sapp RM, Evans WS, Eagan LE, et al. The effects of moderate and high intensity exercise on circulating markers of endothelial integrity and activation in young, healthy men. *J Appl Physiol.* 2019;127:1245-1256.
 132. Oda K, Okada H, Suzuki A, et al. Factors enhancing serum syndecan-1 concentrations: a large-scale comprehensive medical examination. *J Clin Med.* 2019;8(9):1320.
 133. Lee S, Kolset SO, Birkeland KI, Drevon CA, Reine TM. Acute exercise increases syndecan-1 and -4 serum concentrations. *Glycoconj J.* 2019;36:113-125.

134. Swenson KE, Berger MM, Sareban M, et al. Rapid ascent to 4559 m is associated with increased plasma components of the vascular endothelial glycocalyx and may be associated with acute mountain sickness. *High Alt Med Biol.* 2020;21:176-183.
135. Pranskuniene Z, Belousoviene E, Baranauskiene N, et al. Modulation of endothelial glycocalyx and microcirculation in healthy young men during high-intensity sprint interval cycling-exercise by supplementation with pomegranate extract. A randomized controlled trial. *Int J Environ Res Public Health.* 2020;17:4405.
136. Nieuwdorp M, van Haften TW, Gouverneur MC, et al. Loss of endothelial glycocalyx during acute hyperglycemia coincides with endothelial dysfunction and coagulation activation in vivo. *Diabetes.* 2006;55:480-486.
137. Majerczak J, Grandys M, Frołow M, et al. Age-dependent impairment in endothelial function and arterial stiffness in former high class male athletes is no different to that in men with no history of physical training. *J Am Heart Assoc.* 2019;8:e012670. <https://doi.org/10.1161/JAHA.119.012670>
138. Nieuwdorp M, Meuwese MC, Mooij HL, et al. Tumor necrosis factor- α inhibition protects against endotoxin-induced endothelial glycocalyx perturbation. *Atherosclerosis.* 2009;202:296-303.
139. Feurecker M, van Oosterhout WPJ, Feurecker B, et al. Headache under simulated microgravity is related to endocrine, fluid distribution, and tight junction changes. *Pain.* 2016;157:1072-1078.
140. Feurecker M, Feurecker B, Matzel S, et al. Five days of head-down-tilt bed rest induces noninflammatory shedding of L-selectin. *J Appl Physiol.* 2013;115:235-242.
141. Majerczak J, Duda K, Chlopicki S, et al. Endothelial glycocalyx integrity is preserved in young, healthy men during a single bout of strenuous physical exercise. *Physiol Res.* 2016;65:281-291.
142. Hulde N, Rogenhofer N, Brettner F, et al. The CYCLOCALYX study: Ovulatory cycle affects circulating compartments of the endothelial glycocalyx in blood. *Am J Reprod Immunol.* 2018;79(1):e12767. <https://doi.org/10.1111/aji.12767>
143. Nguyen TH, Liu S, Ong GJ, Stafford I, Frenneaux MP, Horowitz JD. Glycocalyx shedding is markedly increased during the acute phase of Takotsubo cardiomyopathy. *Int J Cardiol.* 2017;243:296-299.
144. Neves FM, Meneses GC, Sousa NE, et al. Syndecan-1 in acute decompensated heart failure—association with renal function and mortality. *Circ J.* 2015;79:1511-1519.
145. Ajaero CN, Procter NEK, Chirkov YY, et al. Endothelial dysfunction and glycocalyx shedding in heart failure: insights from patients receiving cardiac resynchronisation therapy. *Heart Vessels.* 2020;35:197-206.
146. Jung C, Fuernau G, Muench P, et al. Impairment of the endothelial glycocalyx in cardiogenic shock and its prognostic relevance. *Shock.* 2015;43:450-455.
147. Nijst P, Cops J, Martens P, et al. Endovascular shedding markers in patients with heart failure with reduced ejection fraction: results from a single-center exploratory study. *Microcirculation.* 2018;25:e12432. <https://doi.org/10.1111/micc.12432>
148. Frydland M, Ostrowski SR, Møller JE, et al. Plasma concentration of biomarkers reflecting endothelial cell- and glycocalyx damage are increased in patients with suspected ST-elevation myocardial infarction complicated by cardiogenic shock. *Shock.* 2018;50:538-544.
149. Bro-Jeppesen J, Johansson PI, Hassager C, et al. Endothelial activation/injury and associations with severity of post-cardiac arrest syndrome and mortality after out-of-hospital cardiac arrest. *Resuscitation.* 2016;107:71-79.
150. Wernly B, Fuernau G, Masyuk M, et al. Syndecan-1 Predicts outcome in patients with ST-segment elevation infarction independent from infarct-related myocardial injury. *Sci Rep.* 2019;9:18367.
151. Solbu MD, Kolset SO, Jenssen TG, et al. Gender differences in the association of syndecan-4 with myocardial infarction: The population-based Tromsø Study. *Atherosclerosis.* 2018;278:166-173.
152. Ollila A, Vikatmaa L, Virolainen J, et al. The association of endothelial injury and systemic inflammation with perioperative myocardial infarction. *Ann Clin Biochem.* 2019;56:674-683. <https://doi.org/10.1177/0004563219873357>
153. Grundmann S, Fink K, Rabadzhieva L, et al. Perturbation of the endothelial glycocalyx in post cardiac arrest syndrome. *Resuscitation.* 2012;83:715-720.
154. Miranda CH, de Carvalho BM, Schmidt A, Marin-Neto JA, Pazin-Filho A. Evaluation of the endothelial glycocalyx damage in patients with acute coronary syndrome. *Atherosclerosis.* 2016;247:184-188.
155. Nemoto T, Minami Y, Yamaoka-Tojo M, et al. Endothelial glycocalyx and severity and vulnerability of coronary plaque in patients with coronary artery disease. *Atherosclerosis.* 2020;302:1-7.
156. Kucur M, Karadag B, Isman FK, et al. Plasma hyaluronidase activity as an indicator of atherosclerosis in patients with coronary artery disease. *Bratisl Lek Listy.* 2009;110:21-26.
157. Xue XJ, Jiang Y, Chen L, Chen SL. Relationship between the endothelial glycocalyx and the extent of coronary atherosclerosis. *Microcirculation.* 2018;25(8):e12504. <https://doi.org/10.1111/micc.12504>
158. Lehnert P, Johansson PI, Ostrowski SR, et al. Coagulopathy in patients with acute pulmonary embolism: a pilot study of whole blood coagulation and markers of endothelial damage. *Scand J Clin Lab Invest.* 2017;77:19-25.
159. Bogner-Flatz V, Braunstein M, Ocker LE, et al. On-the-scene hyaluronan and syndecan-1 serum concentrations and outcome after cardiac arrest and resuscitation. *Mediators Inflamm.* 2019;2019:1-8. <https://doi.org/10.1155/2019/8071619>
160. Cornelis T, Broers NJH, Titulaer DCLM, et al. Effects of ultrapure hemodialysis and low molecular weight heparin on the endothelial surface layer. *Blood Purif.* 2014;38:203-210.
161. Koch J, Idzerda NMA, Ettema EM, et al. An acute rise of plasma Na⁽⁺⁾ concentration associates with syndecan-1 shedding during hemodialysis. *Am J Physiol Renal Physiol.* 2020;319:F171-F177.
162. Campos NG, Marizeiro DF, Florêncio ACL, et al. Effects of respiratory muscle training on endothelium and oxidative stress biomarkers in hemodialysis patients: a randomized clinical trial. *Respir Med.* 2018;134:103-109.
163. Dane MJC, Khairoun M, Lee DH, et al. Association of kidney function with changes in the endothelial surface layer. *Clin J Am Soc Nephrol.* 2014;9:698-704.
164. Kong X, Du J, Su H, et al. Serum levels of the endothelial glycocalyx constituents and the early failure of forearm autologous arteriovenous fistulas in end-stage renal disease patients: a prospective cohort study. *Int Urol Nephrol.* 2020;52:169-177.
165. Padberg J-S, Wiesinger A, di Marco GS, et al. Damage of the endothelial glycocalyx in chronic kidney disease. *Atherosclerosis.* 2014;234:335-343.
166. Vlahu CA, Lemkes BA, Struijk DG, Koopman MG, Krediet RT, Vink H. Damage of the endothelial glycocalyx in dialysis patients. *J Am Soc Nephrol.* 2012;23:1900-1908.
167. Salmato FT, de Oliveira Neves FM, Meneses GC, de Almeida LR, Martins AM, Liborio AB. Glycocalyx injury in adults with nephrotic syndrome: association with endothelial function. *Clin Chim Acta.* 2015;447:55-58.
168. Chaves M, Mendes M, Schwermann M, et al. Angiopietin-2: a potential mediator of the glycocalyx injury in adult nephrotic patients. *J Clin Med.* 2018;7:401.
169. Inkien N, Pettila V, Lakkisto P, et al. Association of endothelial and glycocalyx injury biomarkers with fluid administration, development of acute kidney injury, and 90-day mortality: data from the FINNAKI observational study. *Ann Intensive Care.* 2019;9:103.
170. Hahn RG, Hasselgren E, Bjorne H, Zdosek M, Zdosek J. Biomarkers of endothelial injury in plasma are dependent on kidney function. *Clin Hemorheol Microcirc.* 2019;72:161-168.

171. Li J, Du H, Bai XF, et al. Study on expression of plasma sCD138 in patients with hemorrhagic fever with renal syndrome. *BMC Infect Dis.* 2018;18:100.
172. Meneses GC, Cavalcante MG, da Silva Junior GB, et al. Endothelial glycocalyx damage and renal dysfunction in HIV patients receiving combined antiretroviral therapy. *AIDS Res Hum Retroviruses.* 2017;33:703-710.
173. Meneses GC, Daher EDF, Seguro AC, et al. Endothelial glycocalyx damage is associated with leptospirosis acute kidney injury. *Am J Trop Med Hyg.* 2015;92:611-616.
174. Connolly-Andersen AM, Thunberg T, Ahlm C. Endothelial activation and repair during hantavirus infection: association with disease outcome. *Open Forum Infect Dis.* 2014;29:ofu027.
175. Yeo TW, Weinberg JB, Lampah DA, et al. Glycocalyx breakdown is associated with severe disease and fatal outcome in Plasmodium falciparum malaria. *Clin Infect Dis.* 2019;69:1712-1720.
176. Tang T-C, Alonso S, Ng L-P, et al. Increased serum hyaluronic acid and heparan sulfate in dengue fever: association with plasma leakage and disease severity. *Sci Rep.* 2017;7:46191.
177. Lyimo E, Haslund LE, Ramsing T, et al. In vivo imaging of the buccal mucosa shows loss of the endothelial glycocalyx and perivascular hemorrhages in pediatric Plasmodium falciparum malaria. *Infect Immun.* 2020;88:e00679-e719.
178. Suwanto S, Sasmono RT, Sinto R, Ibrahim E, Suryamin M. Association of endothelial glycocalyx and tight and adherens junctions with severity of plasma leakage in dengue infection. *J Infect Dis.* 2017;251:992-999.
179. Yeo TW, Bush PA, Chen Y, et al. Glycocalyx breakdown is increased in African children with cerebral and uncomplicated falciparum malaria. *FASEB J.* 2019;33:14185-14193.
180. Yeo TW, Weinberg JB, Lampah DA, et al. Glycocalyx breakdown is associated with severe disease and fatal outcome in plasmodium falciparum malaria. *Clin Infect Dis.* 2019;69:1712-1720.
181. Larsen AM, Leino EB, Johansson PI, Birgens H, Ostrowski SR. High syndecan-1 levels in acute myeloid leukemia are associated with bleeding, thrombocytopenia, endothelial cell damage, and leukocytosis. *Leuk Res.* 2013;37:777-783.
182. Kim KJ, Kim JY, Baek IW, Kim WU, Cho CS. Elevated serum levels of syndecan-1 are associated with renal involvement in patients with systemic lupus erythematosus. *J Rheumatol.* 2015;42:202-209.
183. Luo L, Feng S, Wu Y, Su Y, Jing F, Yi Q. Serum levels of syndecan-1 in patients with Kawasaki disease. *Pediatr Infect Dis J.* 2019;38:89-94.
184. Ohnishi Y, Yasudo H, Suzuki Y, et al. Circulating endothelial glycocalyx components as a predictive marker of coronary artery lesions in Kawasaki disease. *Int J Cardiol.* 2019;292:236-240.
185. Nagga K, Hansson O, van Westen D, Minthon L, Wennstrom M. Increased levels of hyaluronic acid in cerebrospinal fluid in patients with vascular dementia. *J Alzheimers Dis.* 2014;42:1435-1441.
186. Larsen AM, Leino EB, Johansson PI, Birgens H, Ostrowski SR. Haemostatic function and biomarkers of endothelial damage before and after RBC transfusion in patients with haematologic disease. *Vox Sang.* 2015;109:52-61.
187. Baron-Stefaniak J, Leitner GC, Kuntzel NKI, et al. Transfusion of standard-issue packed red blood cells induces pulmonary vasoconstriction in critically ill patients after cardiac surgery. A randomized, double-blinded, clinical trial. *PLoS One.* 2019;14(3):e0213000. <https://doi.org/10.1371/journal.pone.0213000>
188. Benediktsson S, Kander T, Ostrowski SR, Johansson PI, Thomas OD, Schott U. Platelet increment is not associated with endothelial damage in haematological patients: a prospective observational study. *Scand J Clin Lab Invest.* 2019;79:395-403.
189. Donati A, Damiani E, Luchetti M, et al. Microcirculatory effects of the transfusion of leukodepleted or non-leukodepleted red blood cells in patients with sepsis: a pilot study. *Crit Care.* 2014;18:R33.
190. Shalaby S, Simioni P, Campello E, et al. Endothelial damage of the portal vein is associated with heparin-like effect in advanced stages of cirrhosis. *Thromb Haemost.* 2020;120:1173-1181.
191. Rovas A, Lukasz AH, Vink H, et al. Bedside analysis of the sublingual microvascular glycocalyx in the emergency room and intensive care unit - the GlycoNurse study. *Scand J Trauma Resusc Emerg Med.* 2018;26:16.
192. Damiani E, Adrario E, Luchetti MM, et al. Plasma free hemoglobin and microcirculatory response to fresh or old blood transfusions in sepsis. *PLoS One.* 2015;10(5):e0122655. <https://doi.org/10.1371/journal.pone.0122655>
193. Donati A, Damiani E, Domizi R, et al. Alteration of the sublingual microvascular glycocalyx in critically ill patients. *Microvasc Res.* 2013;90:86-89.
194. Donati A, Damiani E, Botticelli L, et al. The aPC treatment improves microcirculation in severe sepsis/septic shock syndrome. *BMC Anesthesiol.* 2013;13:25.
195. Rovas A, Seidel LM, Vink H, et al. Association of sublingual microcirculation parameters and endothelial glycocalyx dimensions in resuscitated sepsis. *Crit Care.* 2019;23:260.
196. Beurskens DM, Bol ME, Delhaas T, et al. Decreased endothelial glycocalyx thickness is an early predictor of mortality in sepsis. *Anaesth Intensive Care.* 2020;48:221-228.
197. Drost CC, Rovas A, Kusche-Vihrog K, et al. Tie2 activation promotes protection and reconstitution of the endothelial glycocalyx in human sepsis. *Thromb Haemost.* 2019;119:1827-1838.
198. Martens RJ, Vink H, van Oostenbrugge RJ, Staals J. Sublingual microvascular glycocalyx dimensions in lacunar stroke patients. *Cerebrovasc Dis.* 2013;35:451-454.
199. Ikonomidis I, Frogoudaki A, Vrettou A-R, et al. Impaired arterial elastic properties and endothelial glycocalyx in patients with embolic stroke of undetermined source. *Thromb Haemost.* 2019;119:1860-1868.
200. Ikonomidis I, Pavlidis G, Thymis J, et al. Effects of glucagon-like peptide-1 receptor agonists, sodium-glucose cotransporter-2 inhibitors, and their combination on endothelial glycocalyx, arterial function, and myocardial work index in patients with Type 2 diabetes mellitus after 12-month treatment. *J Am Heart Assoc.* 2020;9:e015716.
201. Ikonomidis I, Pavlidis G, Lambadiari V, et al. Early detection of left ventricular dysfunction in first-degree relatives of diabetic patients by myocardial deformation imaging: the role of endothelial glycocalyx damage. *Int J Cardiol.* 2017;233:105-112.
202. Groen BB, Hamer HM, Snijders T, et al. Skeletal muscle capillary density and microvascular function are compromised with aging and type 2 diabetes. *J Appl Physiol.* 2014;116:998-1005.
203. Wadowski PP, Kautzky-Willer A, Gremmel T, et al. Sublingual microvasculature in diabetic patients. *Microvasc Res.* 2020;129:103971.
204. Nussbaum C, Cavalcanti Fernandes Heringa A, Mormanova Z, Puchwein-Schwepecke AF, Bechtold-Dalla Pozza S, Genzel-Boroviczeny O. Early microvascular changes with loss of the glycocalyx in children with type 1 diabetes. *J Pediatr.* 2014;164:584-589.
205. Wernly B, Fuernau G, Masyuk M, et al. syndecan-1 predicts outcome in patients with ST-segment elevation infarction independent from infarct-related myocardial injury. *Sci Rep.* 2019;2019(9):18367.
206. Wu Q, Gao W, Zhou J, et al. Correlation between acute degradation of the endothelial glycocalyx and microcirculation dysfunction during cardiopulmonary bypass in cardiac surgery. *Microvasc Res.* 2019;124:37-42.
207. Koning NJ, Vonk AB, Vink H, Boer C. Side-by-side alterations in glycocalyx thickness and perfused microvascular density during acute microcirculatory alterations in cardiac surgery. *Microcirculation.* 2016;23:69-74.

208. Nussbaum C, Haberer A, Tiefenthaller A, et al. Perturbation of the microvascular glycocalyx and perfusion in infants after cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2015;150:1474-1481.
209. Astapenko D, Pouska J, Benes J, et al. Neuraxial anesthesia is less harmful to the endothelial glycocalyx during elective joint surgery compared to general anesthesia. *Clin Hemorheol Microcirc.* 2018;72:11-21.
210. Brettner F, Heitzer M, Thiele F, et al. Non-invasive evaluation of macro- and microhemodynamic changes during induction of general anesthesia. A prospective observational single-blinded trial. *Clin Hemorheol Microcirc.* 2020;77(1):1-16. <https://doi.org/10.3233/CH-190691>
211. Haeren R, Rijkers K, Schijns O, et al. In vivo assessment of the human cerebral microcirculation and its glycocalyx: A technical report. *J Neurosci Methods.* 2018;303:114-125.
212. Pouska J, Tegl V, Astapenko D, Cerny V, Lehmann C, Benes J. Impact of intravenous fluid challenge infusion time on macrocirculation and endothelial glycocalyx in surgical and critically ill patients. *Biomed Res Int.* 2018;2018:1-11. <https://doi.org/10.1155/2018/8925345>
213. Nieuwdorp M, Meuwese MC, Mooij HL, et al. Measuring endothelial glycocalyx dimensions in humans: a potential novel tool to monitor vascular vulnerability. *J Appl Physiol.* 2008;104:845-852.
214. Townsend D, D'Aiuto F, Deanfield J. Vascular fragility and the endothelial glycocalyx in the tissues lining the healthy gingival crevice. *J Periodontol.* 2016;87:672-679.
215. Ikonomidis I, Marinou M, Vlastos D, et al. Effects of varenicline and nicotine replacement therapy on arterial elasticity, endothelial glycocalyx and oxidative stress during a 3-month smoking cessation program. *Atherosclerosis.* 2017;262:123-130.
216. Grudyano AI, Tkacheva ON, Khatagov AT, Mustafina FN, Gorshkov AY. The impact of periodontal disease treatment on endothelium of sublingual microvessels. *Stomatologiia (Mosk).* 2016;95:9-12.
217. Schmitz B, Niehues H, Lenders M, et al. Effects of high-intensity interval training on microvascular glycocalyx and associated microRNAs. *Am J Physiol Heart Circ Physiol.* 2019;316:H1538-H1551.
218. Rorije N, Engberink RO, van den Born B, Verberne H, Vogt L. Effects of an acute and chronic salt load on microvascular permeability in healthy subjects (Abstract). *J Hypertens.* 2015; <https://doi.org/10.1097/01.hjh.0000467623.14967.fc>
219. Lee DH, Dane MJC, van den Berg BM, et al. Deeper penetration of erythrocytes into the endothelial glycocalyx is associated with impaired microvascular perfusion. *PLoS One.* 2014;9(5):e96477. <https://doi.org/10.1371/journal.pone.0096477>
220. Wadowski PP, Hülsmann M, Schörgenhofer C, et al. Sublingual functional capillary rarefaction in chronic heart failure. *Eur J Clin Invest.* 2018;48(2):e12869. <https://doi.org/10.1111/eci.12869>
221. Wadowski PP, Steinlechner B, Zimpfer D, et al. Functional capillary impairment in patients with ventricular assist devices. *Sci Rep.* 2019;9:5909.
222. Ikonomidis I, Voumvourakis A, Makavos G, et al. Association of impaired endothelial glycocalyx with arterial stiffness, coronary microcirculatory dysfunction, and abnormal myocardial deformation in untreated hypertensives. *J Clin Hypertens (Greenwich).* 2018;20:672-679.
223. Triantafyllidi H, Benas D, Vlachos S, et al. HDL cholesterol levels and endothelial glycocalyx integrity in treated hypertensive patients. *J Clin Hypertens (Greenwich).* 2018;20:1615-1623.
224. Triantafyllidi H, Benas D, Schoinas A, et al. Sex-related associations of high-density lipoprotein cholesterol with aortic stiffness and endothelial glycocalyx integrity in treated hypertensive patients. *J Clin Hypertens (Greenwich).* 2020;22:1827-1834.
225. Jaarsma C, Vink H, van Haare J, et al. Non-invasive assessment of microvascular dysfunction in patients with microvascular angina. *Int J Cardiol.* 2017;248:433-439.
226. Mulders TA, Nieuwdorp M, Stroes ES, Vink H, Pinto-Sietsma SJ. Non-invasive assessment of microvascular dysfunction in families with premature coronary artery disease. *Int J Cardiol.* 2013;168:5026-5028.
227. Brands J, Hubel CA, Althouse A, Reis SE, Pacella JJ. Noninvasive sublingual microvascular imaging reveals sex-specific reduction in glycocalyx barrier properties in patients with coronary artery disease. *Physiol Rep.* 2020;8:e14351.
228. Amraoui F, Olde Engberink RH, van Gorp J, Ramdani A, Vogt L, van den Born BJ. Microvascular glycocalyx dimension estimated by automated SDF imaging is not related to cardiovascular disease. *Microcirculation.* 2014;21(6):499-505. <https://doi.org/10.1111/micc.12125>
229. Triantafyllidi H, Apostolopoulou O, Benas D, et al. Systemic endothelial glycocalyx and aortic stiffness are preserved in pulmonary arterial hypertension. *Hellenic J Cardiol.* 2019;61(4):288-290. <https://doi.org/10.1016/j.hjc.2019.10.006>
230. Kliche K, Gerth U, Pavenstädt H, Oberleithner H. Recharging red blood cell surface by hemodialysis. *Cell Physiol Biochem.* 2015;35:1107-1115.
231. Meyring-Wösten A, Kuntsevich V, Campos I, et al. Erythrocyte sodium sensitivity and eryptosis in chronic hemodialysis patients. *Kidney Blood Press Res.* 2017;42:314-326.
232. Mitsides N, Cornelis T, Broers NJH, et al. Extracellular overhydration linked with endothelial dysfunction in the context of inflammation in haemodialysis dependent chronic kidney disease. *PLoS One.* 2017;12(8):e0183281. <https://doi.org/10.1371/journal.pone.0183281>
233. Woodford J, Yeo TW, Piera KA, et al. Early endothelial activation precedes glycocalyx degradation and microvascular dysfunction in experimentally induced Plasmodium falciparum and Plasmodium vivax infection. *Infect Immun.* 2020;88:e000895-e919.
234. Miranda S, Billoir P, Le Besnerais M, et al. New insights into antiphospholipid-related endothelial dysfunction by assessment of vascular glycocalyx layer: results from a preliminary cross-sectional study. *Lupus.* 2020;29:157-164.
235. Nelson A, Berkestedt I, Bodelsson M. Circulating glycosaminoglycan species in septic shock. *Acta Anaesthesiol Scand.* 2014;58:36-43.
236. Hahn RG. Renal fluid conservation and the volume kinetics of fluid-induced diuresis. *Clin Exp Pharm Physiol.* 2020; <https://doi.org/10.1111/1440-1681.13439>
237. Chappell D, Jacob M. Role of glycocalyx on fluid management: small things matter. *Best Pract Res Clin Anaesthesiol.* 2014;28:227-234.
238. Rehm M, Haller M, Orth V, et al. Changes in blood volume and hematocrit during acute perioperative volume loading with 5% albumin or 6% hetastarch solutions in patients before radical hysterectomy. *Anesthesiol.* 2001;95:849-856.
239. Jacob M, Chappell D, Rehm M. Clinical update: perioperative fluid management. *Lancet.* 2007;369:1984-1986.
240. Chappell D, Jacob M, Hofmann-Kiefer K, Conzen P, Rehm M. A rational approach to perioperative fluid management. *Anesthesiol.* 2008;109:723-740.
241. Potter DR, Jiang J, Damiano ER. The recovery time course of the endothelial cell glycocalyx in vivo and its implications in vitro. *Circ Res.* 2009;104:1318-1325.
242. Yang Y, Haeger SM, Suflita MA, et al. Fibroblast growth factor signaling mediates pulmonary endothelial glycocalyx reconstruction. *Am J Respir Cell Mol Biol.* 2017;56:727-737.
243. Gabrielsson J, Weiner D. *Pharmacokinetic and Pharmacodynamic Data Analysis*, 4th edn. Swedish Pharmaceutical Society; 2000:193-224.
244. The Human Protein Atlas, sdc-1 and heparan sulfate, available <http://proteinatlas.org>, assessed on December 22, 2020
245. Edul VS, Ince C, Vazquez AR, et al. Similar microcirculatory alterations in patients with normodynamic and hyperdynamic septic shock. *Ann Am Thorac Soc.* 2016;13:240-247.
246. Adamson RH. Permeability of frog mesenteric capillaries after partial pronase digestion of the endothelial glycocalyx. *J*

- Physiol.* 1990;428(1):1-13. <https://doi.org/10.1113/jphysiol.1990.sp018197>
247. Guerri P, Ergin B, Uz Z, et al. Glycocalyx degradation is independent of vascular barrier permeability increase in nontraumatic hemorrhagic shock in rats. *Anesth Analg.* 2019;129:598-607.
 248. Ergin B, Guerri P, Uz Z, et al. Hemodilution causes glycocalyx shedding without affecting vascular endothelial barrier permeability in rats. *J Clin Transl Res.* 2020;5:243-252.
 249. Savary MD, Jiang JX, Park PW, Damiano ER. The endothelial glycocalyx in syndecan-1 deficient mice. *Microvasc Res.* 2013;87:83-91.
 250. Voyvodic PL, Min D, Liu R, et al. Loss of syndecan-1 induces a pro-inflammatory phenotype in endothelial cells with a dys-regulated response to atheroprotective flow. *J Biol Chem.* 2014;289:9547-9559.
 251. Langford JK, Stanley MJ, Cao D, Sanderson RD. Multiple heparan sulfate chains are required for optimal syndecan-1 function. *J Biol Chem.* 1998;273:29965-29971.
 252. Groner W, Winkelmann JW, Harris AG, et al. Orthogonal polarization spectral imaging: a new method for the study of the microcirculation. *Nat Med.* 1999;5:1209-1212.
 253. Goedhart PT, Khalilzadeh M, Bezemer R, Merza J, Ince C. Sidestream dark field (SDF) imaging: a novel stroboscopic LED ring-based imaging modality for clinical assessment of the microcirculation. *Opt Express.* 2007;15:1501-1514.
 254. Gilbert-Kawai E, Coppel J, Bountziouka V, Ince C, Martin D, Caudwell Xtreme Everest and Xtreme Everest 2 Research Groups. A comparison of the quality of image acquisition between incident darkfield and sidestream dark field. *BMC Med Imaging.* 2016;16:10.
 255. Oberleithner H. Sodium selective erythrocyte glycocalyx and salt sensitivity in man. *Pflügers Arch.* 2015;467:1319-1325.
 256. Oberleithner H, Wilhemi M. Determination of erythrocyte sodium sensitivity in man. *Pflügers Arch.* 2013;465:1459-1514.

How to cite this article: Hahn RG, Patel V, Dull RO. Human glycocalyx shedding: Systematic review and critical appraisal. *Acta Anaesthesiol Scand.* 2021;65:590-606. <https://doi.org/10.1111/aas.13797>

APPENDIX

The following methods have been used to assess the thickness of the glycocalyx layer in vivo.

Bright-field imaging	Microcirculation is illuminated with visible light, positioned at a long working distance from the tissues and projected at angles between 45 and 90 degrees from the tissue surface. Reflected image is projected through a lens for direct viewing or capture by a camera. Provides relatively low resolution.
Orthogonal polarization spectral (OPS) imaging	Microcirculation is illuminated with polarized green light. Reflected light from the tissue surface is captured and quantified. Emission and reflected light travel through the same tube thus method is sensitive to internal reflectance. Signal processing includes filtering of reflected polarized light to improve visualization of underlying vessels. ²⁵²
Sidestream dark field (SDF) imaging	Developed as a refinement to OPS. Uses light emitting diodes (LEDs; 530 nm) tuned to absorption by hemoglobin. LEDs are arranged concentrically around a center sensing tube and are placed in contact with tissue. Light paths for excitation and emission are different thus reducing internal interference and providing clearer images. ²⁵³
Incidental dark field (IDF) imaging	Similar to SDF, but IDF illuminates tissue using a non-homogenous field. Incident illumination is projected at a very low angle relative to the tissue surface. Increased signal sensitivity, greater field of view and improved optical resolution contributes to enhance image quality. ²⁵⁴
Erythrocyte sodium sensitivity test (ESST)	Based on the principle that the red blood cell glycocalyx and vascular endothelial glycocalyx are in direct contact. Loss of the endothelial glycocalyx increases drag forces on the RBC surface and damages the RBC glycocalyx. This damage can be assessed and quantified by salt-sensitive sedimentation rates of the RBC. Studies have shown a correlation between EC glycocalyx thickness and RBC sedimentation rates. ^{255,256}